Table 23.7 Cancer of the Prostate (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
Age-Adjusted Rates, 2008-2012						
Age at Diagnosis/Death:						
All ages	137.9	130.4	214.5	21.4	19.8	46.3
Under 65	54.7	51.0	97.6	1.7	1.5	4.3
65 and over	713.5	679.1	1,022.4	157.6	146.5	337.0
All ages (IARC world std) <sup>c</sup>	96.0	90.7	153.3	9.3	8.5	21.0
Age-Specific Rates, 2008-2012						
Age at Diagnosis/Death:						
<1	=	=	=	=	=	-
1-4	-	-	_	_	-	-
5-9	-	-	_	_	-	_
10-14	-	-	_	_	-	_
15-19	_	_	_	_	_	_
20-24	_	_	_	_	_	_
25-29	_	_	_	_	_	_
30-34	0.1	_	_	_	_	_
35-39	1.1	0.9	2.8	0.0	_	_
40-44	10.2	8.7	25.0	0.2	0.1	0.4
45-49	45.5	39.1	105.6	0.9	0.7	2.2
50-54	139.5	128.2	262.3	3.3	2.7	8.3
55-59	302.6	283.1	530.8	8.8	7.5	21.9
60-64	518.6	493.4	856.2	20.1	17.3	50.3
65-69	788.1	753.0	1,207.3	40.4	35.4	101.4
70-74	835.4	800.7	1,167.8	77.4	68.8	189.9
75-79	741.2	700.7	1,004.4	138.8	127.1	312.1
80-84	545.2	512.8	745.3	244.2	228.2	533.8
85+	444.2	420.5	666.4	513.5	491.0	975.7

SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

Rates are per 100,000 and are age-adjusted to the IARC world standard population.

Statistic not shown. Rate based on less than 16 cases for the time interval.